

This file has been cleaned of potential threats.

To view the reconstructed contents, please SCROLL DOWN to next page.

Benefit Maximize by Application of Humic Acid and Magnetized Water for Controlling *Meloidogyne javanica* Infected Khella Plants in Relation to Improve growth and Active Constituents

M.S. Abo-Korah¹ and M.M. Moussa²

¹ Economic Entomology and Agricultural Zoology Dept.,

² Horticulture Dept., Fac. of Agric., Menoufia Univ., Shibin El-Kom, Egypt.

ABSTRACT

Pot experiments were carried out for exploration responsiveness to application of five Humic acid (HA) concentration rates (0,500,1000,1500 and 2000 ppm) and with irrigation by magnetized water (MW), solely or combined for controlling *Meloidogyne javanica* infected khella plants (*Ammi visnaga* L.) in relation to its morphological characters; yield parameters and medicinal components through two winter seasons (2012 and 2013) under shield conditions. Generally, all treatments reduced significantly number of root galls; egg-masses; soil, root population density of the *M. javanica* and increased tested plant morphological characters i.e., plant height; number of main branches; fresh weight of whole plant and yield parameters such as, number of umbels; its fresh and dry weights (g/plant), and fruit yield (g/plant). Dual application of both HA and irrigation with MW combined gave better results more than the solely one in controlling of *M. javanica* and all plant characteristics during the tested two winter seasons. Combined treatment of (HA 1500 ppm + MW) gave the highest reduction percentages in *M. javanica* population (71.0 and 71.8%) through the two successive seasons, respectively. In addition, application of the same combined treatment gave a best results of gall index (1.5 and 1.5); egg-masses production (7.0 and 8.0), in the both seasons, respectively, and a better enhancement in Khella morphological characters and its yield parameters. Highest value of khellin % explicitly was obtained by using 1000 ppm HA+MW and 1500 ppm HA +MW during the two successive winter seasons, respectively.

Key words: Control, *Meloidogyne javanica*, Humic acid, magnetized water, *Ammi visnaga*

INTRODUCTION

Khella plants (*Ammi visnaga* L.) is a member of the family Apiaceae (Umbelliferae); native to the Mediterranean and cultivated area in Egypt (Rechinger, 1972). *A. visnaga* is antiasthmatic, diuretic and vasodilator. It is an effective muscle relaxant and has been used for centuries to alleviate the excruciating pain of kidney stones (Chevallier, 1996). The medicinal properties of *A. visnaga* against coronary diseases and bronchial asthma are attributed to its essential oils that includes the substance 'khellin' (Rose and Hulburd, 1992; Satrani *et al.*, 2004).

Meloidogyne javanica is sedentary endoparasite; its species are among the most destructive agricultural pests, and aggressive to a wide range of crops (Elgorban *et al.*, 2014). Root-knot nematode, *M. javanica* is widely distributed on medicinal & ornamental plants and considered of economic importance to

several species of medicinal plants (Park *et al.*, 1998). Information on host status of medicinal plants for *M. javanica* are important for continued sustainable production and to determine appropriate management practices (Park *et al.*, 2005).

Humic acid is an excellent natural and 100% organic way to provide plants and soil with a concentrated dose of essential nutrients, vitamins and trace elements. Humic acid not only fertilizer but also it acts as a bio-stimulant for the plant (Senn, 1991). There is the most importance need for an eco-friendly substitute for nematode management. Humic acid released during the decomposition of raw organic materials are one of many factors contributing to reduce nematode damage (Mcbride *et al.*, 2000). A new strategy for adjusting plant parasitic nematodes is based on the activation of the plant's own defense system through various biotic and abiotic agents such as Humic acid (El-Sherif *et al.*, 2015).

The most important features of humic acid lie on its ability to bind available insoluble metal ions, oxides and hydroxides, slowly releasing them to roots (Russo and Berlyn 1990) and also prevent water and nutrient from losses in light and sandy soils, as well as, increase aeration and water infiltration in clay soils.

When humic acid entered plants at early stages of development, promoted in increase cell division and root development hastening establishment. In addition, humic acid acted as a natural chelator by enhancing the availability of iron thus increased photosynthesis and sugar production translating into increased storage for defense (Nardi *et al.*, 2002) .

The magnetized water is a kind of water treated by the magnetic field or pass through a magnetic device which causes a positive changes in physical and chemical properties of the water such as pH, electrical conductivity and interfacial tension and so lead to increase of the water quality (Samadyar *et al.*, 2014). Irrigation with the magnetized water increase nutrient mobility in soil and uptake of N, P, K and Fe by plants (Hajer, *et al.*, 2006). It is also increase soil water absorption, and maximize the efficiency of fertilizers. All of these reasons lead to decrease population density of root-knot nematode and increase vegetative growth characters & yield parameters (Koizumi and Kano 2014). Because there is no considerable work hitherto all-over the world generally, and a great scarceness studies herein in Egypt, especially on the effective impact of humic acid and magnetized water and its utility for controlling *M. javanica* infected Khella plants *A. visnaga* in relation to its plant growth properties; yield parameters and medicine components, an attempt had been done to throw light explicitly on this beneficial dual combined appliance of both humic acid and magnetized water and its effectiveness to take into root-knot nematodes control.

MATERIALS AND METHODS

Isolation, Purification and Multiplication

Pot experiments were carried out under shield conditions at the Experimental Farm, Faculty of Agriculture, Minoufia Univ., Shebin El-Kom, Egypt, during two successive winter seasons (2012&2013). Untraditional investigation aimed to scouting the fitness impact of five humic acid (HA) concentration rates (0,500,1000,1500 and 2000 ppm) and irrigation by magnetized water (MW) solely and combinedly for controlling *Meloidogyne javanica* infected khella plants in relation to its growth characters; yield parameters and medicinal components.

Nematode culture

Juveniles of the root-knot nematode, *M. javanica* were obtained from the pure culture reared on black nightshade, planted with *Solanum nigrum* in the Nematode Laboratory of the Entomology and Zoology Department, Faculty of Agriculture , Menoufia University.

Experimental preparation and design

Seeds of khella plants were obtained from Research Center of Medicinal and Aromatic plant Section, Giza, Egypt. The seeds were sown on the middle of October in each growing season in plastic pots 30 cm in diam. Each pot was filled with 4.0 kg of air-dried clay soil. Physical and chemical properties of the soil used in this study were determined according to the method which described by (Cottenie *et al.*,1982), as follows: PH 7.9, Ec 1.73 ds/m; 2.80 % organic matter, 44.24% silt, 3.84% coarse sand, 27.40% fine sand, 23.20% clay, the texture grade was a clay loam soil. The available macronutrients were 2.30% CaCo₃, 0.12% N and 0.25% P₂O₅. Humic acid treatments were prepared and mixed with the experimental soil with five concentration rates (0,500,1000,1500 and 2000 ppm). After sowing the seeds in pots, directly irrigated either with tap water or with magnetically treated water, which passed through magnetron tube of 2 inches in diameter and 4000 Gaus strength (Ameen and Kassim 2009). Fifteen days later, 1000 J₂ of *M. javanica* per 1kg soil were added by pipette into three holes around the base stem of each seedling.

Soil moisture level was kept at about 60% from its water capacity during the growing period by daily weighting. After one and half month from sowing date, plants were thinned to one plant/pot. All pots were received a recommended dose of NPK fertilizers, namely 2 g calcium super phosphate (15.5% P_2O_5), 2 g potassium sulphate (48% K_2O), which was added immediately after thinning, and 4 g ammonium nitrate (33.5% N), which was divided into two equal portions: the first was added immediately after thinning and the second after one and half month later. The experiment layout was randomized complete block design. Ten treatments were applied, each treatment was replicated three times. Khella plants were harvested at the first of May during maturity stage through two successive winter seasons (2012&2013).

Obtained data showed all measurements of khella plants such as: plant height (cm), number of main branches, fresh weight of whole plant (g/plant), number of umbels/plant, fresh and dry weights of umbels (g/plant) and fruit yield (g/plant).

Nematode Extraction and Enumeration:

Each composite soil sample was carefully mixed, and an aliquot of 100 cm³ was processed for nematode extraction according to methods described by (Southey, 1970), each treatment was replicated three times. An aliquant of 1 ml each of nematode suspensions were pipetted off, placed in a Hawksley counting slide and examined by using a stereomicroscope.

Nematode counts were done after 1,2,3,4,5 and 6 months of application, and the identification to generic level were based on morphology of the adult and larval forms, according to the description of (Mai and Lyon, 1975). Roots were carefully washed, and the nematode galls were counted and rated as mentioned in Table (1), as well as one gram per root was stained by acid fuchsin lactophenol to counted root knot nematode stages inside the roots with the aid of a dissecting microscope.

Number of egg-masses root system were assessed by staining the roots with Phloxin-B solution (0.15 g/l tap water) for 20 minutes according to (Daykin and Hussey 1985).

Table (1): Rating scale levels of resistance or susceptible by gall numbers (Southey, 1970)

| Number of galls/ root system | Gall index | Resistance rating |
|------------------------------|------------|------------------------|
| 0 | 0 | Immune |
| 1-2 | 1 | Highly resistant |
| 3-10 | 2 | Resistant |
| 11-30 | 3 | Moderately resistant |
| 31-70 | 4 | Moderately susceptible |
| 71-100 | 5 | Susceptible |
| >100 | 6 | Highly susceptible |

Chemical analysis

Total carbohydrate percentages in the dried herb were determined by using the colorimetric method of (Dubios *et al.*, 1956). Nitrogen, phosphorus and potassium percentages were determined in the dried herb by Kjeldah methods, Spectrophotometricly and Flame photometer methods, respectively, as reported by (Cottenie *et al.*, 1982). Total khellin and visnagin percentages were determined according to the methods described by (Martelli *et al.*, 1984)

Statistical analysis

The obtained data were subjected to analysis of variance (ANOVA) using CoStat Software, Version 6.4 (2008). The mean differences were compared by Least Significant Difference (L.S.D. 5%)

Reduction percentages were computed according to (Abbott formula 1925).

Increase or decrease % = $\frac{\text{Control} - \text{treatment}}{\text{Control}} \times 100$

RESULTS AND DISCUSSION:

In the first season (2012):

Data presented in Table (2) indicated the average numbers of *M. javanica* juveniles per 100 g soil 30, 60, 90, 120, 150 and 180 days after the application of five humic acid (HA) rates (0, 500, 1000, 1500 and 2000 ppm) and with irrigation by magnetized water (MW), solely and combinedly for controlling *M. javanica* infected khella plants under shield conditions.

Statistical analysis indicated that all treatments significantly suppressed the

nematode population in the soil treated after 30, 60, 90, 120, 150 and 180 days in comparison with control treatment.

The highest reduction percentages of the nematode population in the soil, were recorded by application of combined treatment (HA 1500 ppm + MW) followed by (HA 1000 ppm + MW) with (71.0 and 62.0 %), respectively.

The obtained results are in agreement with (Mcbride *et al.*, 2000) who reported that the organic acids released during the decomposition of organic materials are one of many factors contributing to reductions in nematode damage

Table (2) : Impact of humic acid and magnetized water separately and in combination on the population density of root-knot nematode, *Meloidogyne javanica* infected Khella plants and reduction % ,under shield conditions (2012).

| Treatments | Aver. no. of <i>Meloidogyne javanica</i> juveniles/ 100 g of soil | | | | | | |
|----------------------------|---|----------|----------|----------|----------|----------|--------------|
| | Days post-treatments | | | | | | |
| | 30 Days | 60 Days | 90 Days | 120 Days | 150 Days | 180 Days | overall mean |
| Humic acid 500 ppm | 854.3 c | 786.5 c | 712.0 c | 650.1 c | 594.0 c | 506.3 c | 683.9 c |
| Humic acid 1000 ppm | 809.3 d | 778.0 c | 697.0 d | 589.3 e | 510.0 e | 479.9 d | 643.9 d |
| Humic acid 1500 ppm | 757.0 f | 701.3 f | 589.0 h | 512.3 g | 467.5 g | 406.0 f | 572.2 f |
| Humic acid 2000 ppm | 791.0 e | 735.9 d | 625.0 f | 543.1 f | 487.0 f | 438.0 e | 603.3 e |
| magnetized water | 910.0 b | 894.3 b | 839.0 b | 791.0 b | 737.3 b | 704.9 b | 812.7 b |
| H. acid 500 ppm+ m. water | 801.0 d | 719.3 e | 678.5 e | 603.0 d | 560.0 d | 483.0 d | 640.8 d |
| H. acid 1000 ppm+ m. water | 725.0 h | 612.0 h | 499.3 i | 408.0 h | 371.2 i | 291.0 h | 484.4 h |
| H. acid 1500 ppm+ m. water | 707.0 i | 534.3 i | 404.0 j | 203.3 i | 113.4 j | 81.3 i | 340.6 i |
| H. acid 2000 ppm+ m. water | 743.0 g | 671.0 g | 607.0 g | 520.0 g | 394.0 h | 301.0 g | 539.3 g |
| Control | 969.8 a | 1149.1 a | 1320.0 a | 1465.8 a | 1873.0 a | 2019.9 a | 1466.3 a |
| Reduction percentages (%) | | | | | | | |
| Humic acid 500 ppm | 11.9 | 31.6 | 46.1 | 55.7 | 68.3 | 74.9 | 48.1 |
| Humic acid 1000 ppm | 16.6 | 32.3 | 47.2 | 59.8 | 72.7 | 76.2 | 50.8 |
| Humic acid 1500 ppm | 21.9 | 38.9 | 55.4 | 65.1 | 75.0 | 79.9 | 56.0 |
| Humic acid 2000 ppm | 18.4 | 35.9 | 52.6 | 62.9 | 74.0 | 78.3 | 53.7 |
| magnetized water | 6.2 | 22.2 | 36.4 | 46.0 | 60.6 | 65.1 | 39.4 |
| H. acid 500 ppm+ m. water | 17.4 | 37.4 | 48.6 | 58.9 | 70.1 | 76.1 | 51.4 |
| H. acid 1000 ppm+ m. water | 25.2 | 46.7 | 62.2 | 72.1 | 80.2 | 85.6 | 62.0 |
| H. acid 1500 ppm+ m. water | 27.1 | 53.5 | 69.4 | 86.1 | 93.9 | 95.9 | 71.0 |
| H. acid 2000 ppm+ m. water | 23.4 | 41.6 | 54.0 | 64.6 | 78.9 | 85.1 | 53.4 |

Means in each column followed by the same letter (s) are not significantly different at 5% level.

An alternative to pesticide application is that, it may be possible to utilize a scheme of inducible plant defenses (El-Sherif *et al.*, 2015).

In the second season (2013)

Data presented in Table (3) showed that the side effect of applied treatments on the population density of *M. javanica* infected khella plants under shield plantation conditions.

Statistical analysis indicated that all treatments significantly suppressed nematode population in the soil treated

after 30, 60, 90, 120, 150 and 180 days in comparison with control treatment. The combined treatment of (HA 1500 ppm + MW) recorded the least mean of population density of *M. javanica* (352.7 ind's /100g) in the six months.

Highest reduction percentages of the nematode in the soil were recorded in the combined treatments of (HA 1500 ppm + MW) followed by (HA 2000 ppm + MW) with (71.8&61.5), respectively.

The role of humic acid in overcoming the harmful effects of rust diseases in *Faba bean* plant may be due to the increase in

chitinase activity (Abd El-Kareem, 2007). Both HA and MW may play a considerable role in plant growth stimulation and caused promoting for plant natural resistance against plant-parasites such as root-knot nematode and other pests.

Data presented in Table (4) indicated that all treatments decreased the gall index and egg-masses production. The combined treatment of (HA 1500 ppm + MW) recorded the highest reduction percentage in root gall index (-75.0 & -75.0 %) and egg-masses production (-85.7 and -85.2 %) in the two seasons, respectively (2012 & 2013).

Properness; performable and beneficial application of both HA and irrigation by MW by appliance duality combined treatments gave great utility of nematodes control, which is safety and inexpensive in comparison with commercial nematicides application.

On the other hand, these treatments must be introduced and may be useful in pest management programme against different nematode species infected Khella and different other medicinal & ornamental plants.

Results indicated that application of humic acid and magnetized water each solely for controlling *M. javanica* infect Khella plants gave the lowest reduction percentages of nematodes control. While contrariwise occurred in the effectiveness when it appliance combinedly, resulted the highest reduction percentages (HA1500+MW) that reached (71.0 and 71.8%). Interpretation of such finding may be due to the accumulative effectiveness; behaviorism; own extensive properties and its interaction between them, for each of HA and MW in the case of its application duality combined treatments for controlling root knot nematode *M. javanica*.

Table (3) : Impact of humic acid and magnetized water separately and in combination on the population density of root-knot nematode, *Meloidogyne javanica* infected Khella plants and reduction %, under shield conditions (2013).

| Treatments | Aver. no. of <i>Meloidogyne javanica</i> juveniles/ 100 g of soil | | | | | | |
|----------------------------|---|----------|----------|----------|----------|----------|--------------|
| | Days post-treatments | | | | | | |
| | 30 Days | 60 Days | 90 Days | 120 Days | 150 Days | 180 Days | overall mean |
| Humic acid 500 ppm | 871.0 c | 802.0 c | 723.0 c | 597.0 c | 511.3 c | 467.0 c | 661.9 c |
| Humic acid 1000 ppm | 830.0 d | 789.0 d | 701.0 d | 557.0 d | 467.3 d | 401.0 d | 624.2 d |
| Humic acid 1500 ppm | 710.0 g | 661.3 g | 521.0 g | 482.0 f | 400.3 f | 351.0 f | 520.9 g |
| Humic acid 2000 ppm | 731.0 f | 698.0 e | 512.0 h | 473.0 g | 421.0 e | 383.0 e | 536.3 f |
| magnetized water | 934.0 b | 887.0 b | 806.0 b | 761.0 b | 697.0 b | 643.3 b | 788.1 b |
| H. acid 500 ppm+ m. water | 745.0 e | 697.0 e | 604.0 e | 491.3 e | 405.3 f | 347.0 f | 548.3 e |
| H. acid 1000 ppm+ m. water | 711.0 g | 644.9 h | 531.0 f | 397.0 h | 341.0 h | 298.9 h | 487.3 h |
| H. acid 1500 ppm+ m. water | 671.0 h | 527.3 i | 383.9 i | 304.0 i | 167.0 i | 62.9 i | 352.7 i |
| H. acid 2000 ppm+ m. water | 720.0 G | 687.0 f | 600.3 e | 405.0 h | 361.3 g | 309.0 g | 513.8 g |
| Control | 1008.0 a | 1216.0 a | 1401.0 a | 1567.0 a | 1786.0 a | 2184.0 a | 1527.0 a |
| Reduction percentages (%) | | | | | | | |
| Humic acid 500 ppm | 13.6 | 34.0 | 48.4 | 61.9 | 71.4 | 78.6 | 51.3 |
| Humic acid 1000 ppm | 17.7 | 35.1 | 49.9 | 64.5 | 73.8 | 81.6 | 53.8 |
| Humic acid 1500 ppm | 29.7 | 45.6 | 62.8 | 69.2 | 77.6 | 83.9 | 61.4 |
| Humic acid 2000 ppm | 27.5 | 42.6 | 63.5 | 69.8 | 76.4 | 82.5 | 60.4 |
| magnetized water | 7.3 | 27.1 | 42.5 | 51.4 | 60.9 | 70.5 | 43.3 |
| H. acid 500 ppm+ m. water | 26.1 | 42.7 | 56.9 | 68.7 | 77.3 | 84.1 | 59.3 |
| H. acid 1000 ppm+ m. water | 29.5 | 46.9 | 62.1 | 74.7 | 80.9 | 86.3 | 60.4 |
| H. acid 1500 ppm+ m. water | 33.4 | 56.6 | 72.6 | 80.6 | 90.7 | 97.1 | 71.8 |
| H. acid 2000 ppm+ m. water | 28.6 | 43.5 | 57.2 | 74.2 | 79.7 | 85.9 | 61.5 |

Means in each column followed by the same letter (s) are not significantly different at 5% level.

Table (4): Increase or decrease of gall index and egg-masses production on Khella as influenced by treatments application in the two seasons (2012 and 2013).

| Treatments | 1st season | | 2nd season | |
|----------------------------|-----------------|-----------------------|-----------------|-----------------------|
| | Root gall index | Egg-masses production | Root gall index | Egg-masses production |
| Humic acid 500 ppm | 5.0 ab | 41.0 b | 4.5 b | 36.0 b |
| Humic acid 1000 ppm | 4.0 bc | 31.0 c | 3.5 bcd | 27.0 c |
| Humic acid 1500 ppm | 3.0 cd | 24.0 d | 3.0 cd | 23.0 c |
| Humic acid 2000 ppm | 4.0 bc | 30.0 c | 3.5 bcd | 26.0 c |
| Magnetized water | 5.0 ab | 38.0 b | 4.0 bc | 34.0 b |
| H. acid 500 ppm+ m. water | 3.0 cd | 22.0 d | 2.5 de | 18.0 d |
| H. acid 1000 ppm+ m. water | 2.0 de | 12.0 e | 2.5 de | 17.0 d |
| H. acid 1500 ppm+ m. water | 1.5 e | 7.0 f | 1.5 e | 8.0 e |
| H. acid 2000 ppm+ m. water | 2.0 de | 13.0 e | 3.0 cd | 19.0 d |
| Control | 6.0 a | 49.0 a | 6.0 a | 54.0 a |
| LSD 5% | 0.9 | 3.4 | 0.9 | 3.4 |
| Increase or decrease (%) | | | | |
| Humic acid 500 ppm | -16.6 | -16.3 | -25.0 | -33.3 |
| Humic acid 1000 ppm | -33.3 | -36.7 | -41.7 | -50.0 |
| Humic acid 1500 ppm | -50.0 | -51.0 | -50.0 | -57.4 |
| Humic acid 2000 ppm | -33.3 | -38.8 | -41.7 | -51.9 |
| magnetized water | -16.6 | -22.0 | -33.3 | -37.0 |
| H. acid 500 ppm+ m. water | -50.0 | -55.1 | -58.3 | -66.7 |
| H. acid 1000 ppm+ m. water | -66.6 | -75.5 | -58.3 | -68.5 |
| H. acid 1500 ppm+ m. water | -75.0 | -85.7 | -75.0 | -85.2 |
| H. acid 2000 ppm+ m. water | -66.6 | -73.5 | -50.0 | -64.9 |

Means in each column followed by the same letter (s) are not significantly different at 5% level.

Morphological Characteristics

After six months from sowing, responsiveness of khella plants growth to the application of different rates of humic acid (HA) was determined by means of measuring height, number of main branches and fresh weight of whole plant. Data in Table (5) markedly indicated that, all HA treatments significantly increased plant growth parameters with various extents. In this respect, application of HA

at 1500 ppm ranked the first. The increments in growth parameters may be due to that HA is extremely important component because it constitutes a stable fraction of carbon, thus regulating the carbon cycle and release of nutrients, including nitrogen, phosphorus, and sulfur. Similar findings obtained by (Mohammadipour *et al.*, 2012) on *Calendula officinalis*.

Table (5): Effect of Humic acid, irrigation with both tap & magnetized water and their interaction on vegetative growth characters of *Ammi visnaga* L. during the two growing winter seasons (2012 and 2013).

| First season (2012) | | | | | | | Second season (2013) | | | | | |
|-----------------------------|-------------|---------------|----------------|----------------|----------------|---------|----------------------|---------------|----------------|----------------|----------------|--------|
| Plant height (cm) | | | | | | | | | | | | |
| Treatments | HA 0 ppm | HA 500 ppm | HA 1000 ppm | HA 1500 ppm | HA 2000 ppm | Mean | HA 0 ppm | HA 500 ppm | HA 1000 ppm | HA 1500 ppm | HA 2000 ppm | Mean |
| TW | 81.74 | 84.39 | 88.67 | 91.06 | 89.75 | 87.12 | 83.62 | 84.93 | 90.60 | 95.23 | 91.43 | 89.16 |
| MW | 86.54 | 91.62 | 97.53 | 105.41 | 102.33 | 96.68 | 89.70 | 95.21 | 100.33 | 115.56 | 104.79 | 101.11 |
| mean | 84.14 | 88.00 | 93.10 | 98.23 | 96.04 | - | 86.66 | 90.07 | 95.46 | 105.39 | 98.11 | - |
| L.S.D.at 5% | W= 4.51 | H= 3.57 | | W*H=4.99 | | W= 5.77 | | H=5.23 | | W*H= 7.32 | | |
| Number of main branches | | | | | | | | | | | | |
| TW | 6.35 | 6.89 | 7.48 | 8.15 | 7.78 | 7.33 | 6.69 | 7.15 | 7.50 | 7.95 | 7.46 | 7.35 |
| MW | 7.10 | 7.41 | 8.00 | 8.65 | 8.52 | 7.93 | 7.45 | 8.12 | 8.69 | 9.29 | 8.84 | 8.47 |
| mean | 6.72 | 7.15 | 7.74 | 8.40 | 8.15 | - | 7.07 | 7.63 | 8.09 | 8.62 | 8.15 | - |
| L.S.D.at 5% | W= NS | H= 1.12 | | W*H= 1.57 | | W= 0.89 | | H=1.1 | | W*H=1.54 | | |
| Fresh weight of whole plant | | | | | | | | | | | | |
| TW | 188.83 | 195.62 | 212.54 | 225.29 | 219.71 | 208.39 | 192.05 | 206.37 | 217.11 | 234.40 | 228.51 | 215.68 |
| MW | 211.81 | 223.54 | 245.07 | 256.92 | 238.43 | 235.15 | 218.15 | 225.35 | 251.70 | 259.92 | 246.67 | 240.35 |
| mean | 200.32 | 209.58 | 228.80 | 241.10 | 229.07 | - | 205.10 | 215.86 | 234.40 | 247.16 | 237.59 | - |
| L.S.D.at 5% | W= 7.87 | H= 5.64 | | W*H=7.89 | | W= 8.61 | | H= 7.25 | | W*H= 10.15 | | |

W= watering H= Humic acid w*H= Interaction

The same data clearly indicated that, watering khella plants with magnetized water (MW) raised growth parameters in comparison with those irrigated with tap water (TW) in the two experimental seasons. These results could be explained through the finding of (Midan and Tantawy 2013) who mentioned that irrigation snap beans with magnetized water increased soil content from available N,P and K and consequently improved its absorption by

roots, which was reflected in promoting vegetative growth characters.

Also, there are significant effects of all interactions treatments between HA and irrigation with MW on vegetative growth parameters. The combined treatment of MW with HA at rate of (1500 ppm) produced the highest mean values of plant height; number of main branches and fresh weight of whole plant during the both seasons.

Table (6): Effects of Humic acid, irrigation with both tap & magnetized water and their interaction on yield parameters of *Ammi visnaga* L. during the two growing winter seasons (2012 and 2013).

| First season (2012) | | | | | | Second season (2013) | | | | | | |
|----------------------------------|----------|------------|-------------|-------------|-------------|----------------------|----------|------------|-------------|-------------|-------------|-------|
| Treatments | HA 0 ppm | HA 500 ppm | HA 1000 ppm | HA 1500 ppm | HA 2000 ppm | Mean | HA 0 ppm | HA 500 ppm | HA 1000 ppm | HA 1500 ppm | HA 2000 ppm | Mean |
| Number of umbels/plant | | | | | | | | | | | | |
| TW | 14.49 | 15.26 | 15.34 | 18.02 | 16.65 | 15.95 | 15.77 | 16.40 | 17.86 | 19.94 | 19.36 | 17.86 |
| MW | 16.56 | 18.08 | 19.17 | 22.41 | 20.37 | 19.31 | 18.22 | 20.81 | 23.65 | 25.73 | 21.32 | 21.94 |
| mean | 15.52 | 16.67 | 17.25 | 20.21 | 18.51 | - | 16.99 | 18.60 | 20.75 | 22.83 | 20.34 | - |
| L.S.D.at 5% | W= 2.58 | | H= 2.04 | | W*H= 2.86 | | W= 3.01 | | H= 2.87 | | W*H=4.02 | |
| Fresh weight of umbels(g/plant) | | | | | | | | | | | | |
| TW | 56.40 | 58.35 | 63.67 | 67.25 | 65.72 | 62.27 | 57.51 | 61.84 | 65.11 | 70.25 | 68.42 | 64.62 |
| MW | 63.33 | 66.92 | 73.58 | 76.89 | 71.40 | 70.42 | 64.37 | 70.53 | 77.29 | 79.75 | 75.82 | 73.55 |
| mean | 59.86 | 62.63 | 68.62 | 72.07 | 68.56 | - | 60.94 | 66.18 | 71.20 | 75.00 | 72.12 | - |
| L.S.D.at 5% | W=6.74 | | H=5.87 | | W*H=8.22 | | W=6.41 | | H=5.99 | | W*H= 8.39 | |
| Dry weight of umbels (g/plant) | | | | | | | | | | | | |
| TW | 33.18 | 35.93 | 39.56 | 41.55 | 40.17 | 38.07 | 37.11 | 38.37 | 43.82 | 45.66 | 44.19 | 41.83 |
| MW | 38.95 | 40.60 | 46.29 | 47.51 | 43.76 | 43.42 | 39.68 | 44.15 | 48.75 | 51.09 | 46.05 | 45.94 |
| mean | 36.06 | 38.26 | 42.92 | 44.53 | 41.96 | - | 38.39 | 41.26 | 46.28 | 48.37 | 45.12 | - |
| L.S.D.at 5% | W= 4.52 | | H=4.34 | | W*H= 6.08 | | W= 2.65 | | H= 2.34 | | W*H=3.28 | |
| Fruit yield (g/plant) | | | | | | | | | | | | |
| TW | 11.86 | 12.95 | 14.35 | 15.05 | 14.78 | 13.79 | 12.67 | 13.37 | 15.13 | 15.80 | 15.41 | 14.47 |
| MW | 14.13 | 14.76 | 16.80 | 17.15 | 15.82 | 15.73 | 13.65 | 15.43 | 16.87 | 17.85 | 16.27 | 16.01 |
| mean | 12.99 | 13.85 | 15.57 | 16.10 | 15.30 | - | 13.16 | 14.40 | 16.00 | 16.82 | 15.84 | - |
| L.S.D.at 5% | W=1.52 | | H=1.13 | | W*H=1.58 | | W=1.16 | | H=1.04 | | W*H=1.45 | |

W= watering H= Humic acid w*H= Interaction

Yield parameters

Concerning the effect of HA, data presented in Table (6) reported that, yield and yield components such as number of umbels/plant, fresh and dry weights of umbels (g/plant) as well as fruit yield (g/plant) were significantly affected by application of different HA rates and reached its maxima by using 1500 ppm HA in the two seasons. These enhancements in the yield components referred to the favorable effectiveness of HA which has many beneficial effects on soil structure; soil microbial populations and increase

modify mechanisms involved in plant growth stimulation, cell permeability and nutrient uptake. The present results are agree with those of (Khalid *et al.*, 2015) on *Foeniculum vulgare*.

In addition, it is evident that, yield and yield components were enhanced by the application of MW as compared with TW in the winter seasons of 2012 and 2013. The availableness of irrigation with MW may be due to the increase in pigments, photosynthetic rate, IAA, and protein biosynthesis (Hozayn and Abdul qados, 2010). Also watering with MW

enhancement surface tension, polarity, conductivity, hydrogen bonding, pH and solubility of salts in soil. These results

could be also supported by (Ameen and Kassim 2009) on *Jerbera jamesonii*

Table (7): Effects of Humic acid, irrigation with both tap & magnetized water and its interaction on chemical composition and active constituents of *Ammi visnaga* L. during the two growing seasons (2012 and 2013).

| First season (2012) | | | | | | Second season (2013) | | | | | | |
|------------------------|----------------|------------------|-------------------|-------------------|-------------------|----------------------|----------------|------------------|-------------------|-------------------|-------------------|-------|
| Treatments | HA 0 ppm | HA 500 ppm | HA 1000 ppm | HA 1500 ppm | HA 2000 ppm | Mean | HA 0 ppm | HA 500 ppm | HA 1000 ppm | HA 1500 ppm | HA 2000 ppm | Mean |
| N (%) | | | | | | | | | | | | |
| TW | 2.34 | 2.87 | 3.21 | 3.14 | 2.94 | 2.90 | 2.15 | 2.56 | 2.91 | 2.72 | 2.70 | 2.60 |
| MW | 2.88 | 3.33 | 3.52 | 3.24 | 3.05 | 3.20 | 2.62 | 2.81 | 3.19 | 3.18 | 2.90 | 2.94 |
| mean | 2.61 | 3.10 | 3.36 | 3.19 | 2.99 | - | 2.38 | 2.68 | 3.05 | 2.95 | 2.80 | - |
| P (%) | | | | | | | | | | | | |
| TW | 0.28 | 0.34 | 0.39 | 0.39 | 0.35 | 0.35 | 0.30 | 0.28 | 0.35 | 0.40 | 0.38 | 0.34 |
| MW | 0.36 | 0.38 | 0.37 | 0.43 | 0.41 | 0.39 | 0.39 | 0.41 | 0.42 | 0.42 | 0.37 | 0.40 |
| mean | 0.32 | 0.36 | 0.38 | 0.41 | 0.38 | - | 0.34 | 0.34 | 0.38 | 0.41 | 0.37 | - |
| K (%) | | | | | | | | | | | | |
| TW | 2.22 | 2.31 | 2.96 | 2.64 | 2.71 | 2.56 | 2.85 | 2.92 | 3.14 | 3.07 | 2.96 | 2.98 |
| MW | 3.12 | 3.35 | 3.41 | 3.18 | 3.17 | 3.24 | 2.92 | 3.30 | 3.73 | 3.58 | 3.62 | 3.43 |
| mean | 2.67 | 2.83 | 3.18 | 2.91 | 2.94 | - | 2.88 | 3.11 | 3.43 | 3.32 | 3.29 | - |
| Total Carbohydrate (%) | | | | | | | | | | | | |
| TW | 12.15 | 12.62 | 13.03 | 13.42 | 13.51 | 12.94 | 12.29 | 12.41 | 12.85 | 13.22 | 13.74 | 12.90 |
| MW | 12.61 | 13.14 | 13.09 | 13.52 | 13.87 | 13.24 | 13.11 | 13.24 | 13.73 | 13.80 | 14.01 | 13.57 |
| mean | 12.38 | 12.88 | 13.06 | 13.47 | 13.69 | - | 12.70 | 12.82 | 13.29 | 13.51 | 13.87 | - |
| Total khellin % | | | | | | | | | | | | |
| TW | 0.98 | 1.04 | 1.17 | 1.11 | 1.03 | 1.06 | 0.87 | 1.10 | 1.15 | 1.22 | 1.16 | 1.10 |
| MW | 1.21 | 1.26 | 1.32 | 1.25 | 1.28 | 1.26 | 1.19 | 1.17 | 1.21 | 1.28 | 1.26 | 1.22 |
| mean | 1.09 | 1.15 | 1.24 | 1.18 | 1.15 | - | 1.03 | 1.13 | 1.18 | 1.25 | 1.21 | - |
| Total visnagin (%) | | | | | | | | | | | | |
| TW | 0.32 | 0.36 | 0.34 | 0.28 | 0.30 | 0.32 | 0.25 | 0.31 | 0.30 | 0.27 | 0.27 | 0.28 |
| MW | 0.23 | 0.21 | 0.25 | 0.20 | 0.21 | 0.22 | 0.21 | 0.24 | 0.26 | 0.26 | 0.20 | 0.23 |
| mean | 0.27 | 0.28 | 0.29 | 0.24 | 0.25 | - | 0.23 | 0.27 | 0.28 | 0.26 | 0.23 | - |

Generally, combined treatments of HA and MW gave enhancement of the yield and yield components, that recorded the highest mean value in dual treatment of MW+1500 ppm HA.

Total khellin and visnagin contents

Data in Table (7) showed that increasing in HA rates till 1000 ppm means increments in khellin and visnagin contents and then decreased gradually by increasing HA rates up to 2000 ppm. Such quality improvements with HA are in agreements with those obtained by (Khalid *et al.*, 2015) on *Foeniculum vulgare* who emphasized that, the application of HA caused an increment in essential oil percentage and its main constituents i.e., trans-Anethole, Limonene and Fenchone.

Treated khella plants with MW revealed progressive increases in total khellin

content. On the other hand, there was a decline in total visnagin content compared with control plants during both seasons. These results are fortified by those of (Ali *et al.*, 2011) on *Capsicum annuum* whom reported that, magnetic treatments led to a remarkable increase in vit.C and capsaicin contents compared with control plants.

Duality appliance of HA rates and MW and their interaction resulted increments in total khellin content in comparison with those obtained by applying HA rates with TW. Combined treatments of MW + 1000 ppm HA; MW + 1500 ppm HA gave the best results in the first and second seasons, respectively. Meanwhile, the visnagin values showed the reverse and reached to the maximum results by appliance of TW + 500 ppm HA in the two successive growing seasons.

Mineral and total Carbohydrate percentages

In dried herb of khella plants, increments in N, P, K and total carbohydrate percentages occurred parallel with increasing HA rates at its appliance. Application of HA at 1000 ppm gave the highest percentages of N and K, while its adequacy at 1500 and 2000 ppm caused the highest percentages of P and total carbohydrate, respectively in the two winter seasons. Similar results are harmony with that obtained results by (Radwan *et al.*, 2015) on wheat.

Solely treatment irrigation by MW produced the highest N, P, K and total carbohydrate percentages compared with that found by application of TW in irrigation Khella plants through both winter seasons under study. These results are in harmony of that found by (Mazrou 2013) on *Carum carvi*.

All combined treatments of HA+MW (interactions) caused increments of mineral and total carbohydrate percentages. Reached its maxima of N and K % in the treatment of HA1000 ppm + MW, while the highest % of P and total carbohydrate occurred by application of both combined treatment HA1500 and 2000 ppm + MW for each, respectively, in both seasons.

Finally, it could be recommended with appliance of the best and optimum combined treatment among Humic acid rates (1500ppm) plus magnetized water for controlling root-knot nematode *M. javanica* infected Khella plants, hence it gave the highest reduction percentages on it. Consequently, this dual treatment caused enhancement and improvement of Khella morphological characteristics; yield parameters; medicinal components such as total khellin and visnagin contents.

REFERENCES

Abbott, W.S. (1925). A method of computing the effectiveness of an insecticide. J. Econ. Entomol.; 18 : 265-267.

Abd El-Kareem, F. (2007). Induced resistance in bean plants against root rot and Alternaria leaf spot diseases

using biotic and abiotic inducers under field conditions. Res.J. Agric. and Biol. Sci., 3(6): 767-774.

Ali,T.B.; Khalil, S.E. and Khalil, A. M. (2011). Magnetic Treatments of *Capsicum annuum* L. grown under saline irrigation conditions. J. Appl. Sci. Res., 7(11): 1558-1568.

Ameen, S.K. and Kassim, A.F. (2009). Influence of magnetized saline irrigating water and vegetative growth of *Gerbera jamesonii* J. Damascus Univ. for Agric. Sci., 25(1): 63-74.

Chevallier, A. (1996). The Encyclopedia of medicinal Plants. Dorling Kindersley, London, ISBN 9-780751-303148.

Cottenie, A.; Verloo, M.; Kiekens, L.; Velghe, G. and Camerlynck, R. (1982). Chemical analysis of plant and soil. PP. Laboratory of Analytical and Agro chemistry, State.Univ. Ghent. Belgium.3;111-116.

Daykin, M. E., and Hussey, R.S. (1985). Staining and histopathological techniques in nematology. In: Barker, K. R.; C. C. Carter and J. N. Sasser (eds), An Advanced treatise on *Meloidogyne*, Vo. II Methodology, pp. 39-48. North Carolina State University Graphics, Raleigh.

Dubois, M.; Gilles, K.A.; Hamilton, J.K.; Robers P.A. and Smith, F. (1956). Calorimetric method for determination of sugars and related substance. Anal. Chem. 28; 350-356.

Elgorban, A. M.; Wahab, M. A.; Bahkali, A. H. and Al-Sum, B. A. (2014). Biocontrol of *Meloidogyne javanica* on Tomato with *Trichoderma harzianum*. Clean – Soil, Air, Water, 42 (10), 1464–1469.

El-Sherif, A.G.; Gad, S. B.; Khalil, A.M. and Mohamedy, H.E. (2015). Impact of Four organic acids on *Meloidogyne incognita* infecting tomato plants under greenhouse conditions. G.J.B.A.H.S., Vol.4(2):94-100.

Hajer, A.S.; Malibari, A.A.; Al-Zahrani, H.S. and Almaghrabi, O.A. (2006). Responses of three tomato cultivars to

- sea water salinity 1. Effect of salinity on the seedling growth. *African Journal of Biotechnology* 5: 855-861.
- Hozayn, M. and Abdul qados, A.M. (2010). Irrigation with magnetized water enhances growth, chemical constituent and yield of chick pea (*Cicer arietinum* L.). *Agric. Biol. J. North America*, 1(4): 671 - 676.
- Khalid, K. A.; Omer, E.A.; El Gendy, A.G. and Hussein, M.S. (2015). Impact of organic compost and humic acid on essential oil composition of sweet fennel (*Foeniculum vulgare* var. Dulce) under sandy soil conditions in Egypt. *World J Pharm Sci.*, 3(2): 160-166.
- Koizumi, M. and Hiromi K. (2014). Water channel for dry broad bean seeds at germination observed by micromagnetic resonance imaging. *Am. J. Biol. Life Sci.*, 2(2): 37-40.
- Mai, W.F., and Lyon, H. H. (1975). Plant-parasitic nematodes: A Pictorial Key to genera (Comstock Books) Publisher: Peter G. Mullin, Cornell University Press; 5 Sub edition, 277 pp.
- Martelli, P.; Bovalimi, L.; Ferri, S. and Franchi, G.G. (1984). Rapid separation and quantitative determination of Khellin and Visnagin in *Amm visnaga* L. Lam. Fruits by high performance liquid chromatography. *J. chromatography* 301: 297-302.
- Mazrou, R.M. (2013). Physiological studies on *Carum carvi* L. plants P.HD. Thesis, Fac. Agric. . Menoufia Univ., Egypt.
- McBride, R.G.; Mikkelsen, R.L. and Barker, K.R. (2000). The role of low molecular weight organic acids from decomposing rye in inhibiting root-knot nematode populations in soil. *APPI Soil Ecol.*;15:243–251.
- Midan, S. A. and Tantawy, M.F. (2013). Effect of magnetized irrigation water on mineral nitrogen fertilization efficiency, nutrients availability and snap bean plants productivity. *Minufiya J. Agric. Res.*, 38 No.2 (1): 423 - 443.
- Mohammadipour, E. A.; Golchin, J.; Mohammadi, N.; and Zarchini, M. (2012). Effect of humic acid on yield and quality of Marigold (*Calendula officinalis* L.) *Annals of Biological Research*, 3 (11):5095-5098.
- Nardi, S.; Pizzeghello, D.; Muscolob, A. and Vianello, A. (2002). Physiological effects of humic substances on higher plants. *Soil Biol. Biochem.*, 34:1527-36.
- Park, S.D.; Khan, Z.; Choi, Y.E. and Choi, B.S. (1998). Association and distribution of plant-parasitic nematodes in peony (*Paeonia lactiflora*) fields of Korea. *Int J Nematol.*, 8:57–60.
- Park, S. D.; Khan, Z.; Ryu, J. G.; Seo, Y. J. and Yoon, J. T. (2005). Effect of initial density of *Meloidogyne hapla* on its pathogenic potential and reproduction in three species of medicinal plants *J. Phytopathology*, 153: 250–253.
- Radwan, F. I.; Gomaa, M. A.; Rehab, I. F. and Adam, S. I. (2015). Impact of humic acid application, foliar Micronutrients and Biofertilization on Growth, Productivity and quality of Wheat (*Triticum aestivum*, L.). *Middle East J. Agric. Res.*, 4(2): 130-140.
- Rechinger, K.H. (1972). Family umbelliferae, In: flora Iranica, Akademische Druck-und. Verlagsanstalt, Graz, Austria.7;35-42.
- Rose, J. and Hulburd, J. (1992). The aromatherapy book applications and inhalations. North Atlantic Books, Berkeley, California, 94:712-719.
- Russo, R.O. and Berlyn, G.P. (1990). The use of organic biostimulants to help low input sustainable agriculture. *J. Sust Agric.*, (1):19-42.
- Samadyar, H.; Rahi, A.R.; Shirmohammadi, K.; Taghizade, F. and Kadkhoda, Z. (2014). The effects of water electronic filtration (magnetic water) on alkaloids hyoscyamine seeds and some morphological traits in two species of *Datura*. *Journal of Plants and Ecosystems*. 10(40): 59-72. (In Farsi).

Satrani, B.; Farah, A.; Fechtal, M.; Talbi, M. and Boumari, M.L.(2004). Chemical composition and antimicrobial and antifungal activities of the essential oil of *Ammi visnaga* (L.) Lam. Acta. Bot. Gal., 151: 65-71.

Senn, T.L. (1991). Humates in Agriculture. Acres USA, Jan.

Southey, J. F. (1970). Laboratory Methods for work with Plant and soil nematodes. ministry of Agriculture, Fishers and Food. Technical Bulletin 2: 5 th ed., 148 pp.

معظم الفائدة باستخدام الحمض الدبالي Humic acid والمياه الممغنطة في مكافحة نيماتودا تعقد الجذور *Meloidogyne javanica* والتي تصيب نبات الخلة وعلاقة ذلك بتحسين نموها ومكوناتها الفعالة.

محمد سعيد أبوقورة¹ & محمد محمد موسى²

¹قسم الحشرات الاقتصادية والحيوان الزراعي & ²قسم البساتين - كلية الزراعة - جامعة المنوفية - مصر

الملخص العربي

أجريت التجارب في أصص لدراسة تأثير استخدام الحمض الدبالي humic acid بخمسة معدلات وهي (صفر & 500 & 1000 & 1500 & 2000 ppm جزء في المليون) والري بالمياه الممغنطة في معاملات منفردة ومركبة لمكافحة نيماتودا تعقد الجذور *Meloidogyne javanica* والتي تصيب نباتات الخلة وعلاقة ذلك بالصفات المورفولوجية & محصول الثمار للنبات والمكونات الفعالة خلال موسمي الشتاء للعامين 2012 & 2013 تحت ظروف المغطاة في مزرعة التجارب بكلية الزراعة - جامعة المنوفية - شبين الكوم - مصر.

بصفة عامة أدت جميع المعاملات إلى خفض معنوي لأعداد كل من العقد النيماتودية & كتل البيض & الكثافة العددية لنيماتودا التعقد في كل من التربة والجذور مما أدى إلى زيادة في كل من الصفات المورفولوجية لنباتات الخلة مثل طول النبات & عدد الأفرع الرئيسية & الوزن الطازج للنبات ككل وكذلك مكونات المحصول مثل عدد النورات الخيمية ووزنها الطازج والجاف (جم/نبات) ومحصول الثمار (جم/نبات).

أدى تطبيق الري بالمياه الممغنطة مع الحمض الدبالي في معاملات ثنائية إلى إعطاء نتائج جيدة وقاطعة إذا ما قورنت باستعمال كل منهما على حدة (بانفراد) في مقاومة نيماتودا تعقد الجذور *M. javanica* وكذلك تحسن في جميع صفات النباتات في موسمي الشتاء خلال عامي الدراسة.

كما أدت المعاملة الثنائية بين الحمض الدبالي بمعدل 1500 + المياه الممغنطة عن أعلى معدل نسبة إنخفاض في أعداد النيماتودا *M. javanica* خلال الموسمين المتتابعين للدراسة (71.0% & 71.8%) على التوالي إضافة إلى ذلك أعطت نفس هذه المعاملة أفضل النتائج فيما يخص العقد الجذرية (1.5 & 1.5) وإنتاج كتل البيض (7.0 & 8.0) في كلا من الموسمين.

كما أدت هذه المعاملة الثنائية إلى تحسن ملموس في الصفات المورفولوجية للخلة ودلالات إنتاج محصولها. كما تم الحصول على أعلى قيمة للمادة الفعالة وهي الخلين عند تطبيق كل من المعاملتين: الهيوميك أسيد 1000 ppm + المياه الممغنطة & الهيوميك أسيد 1500 ppm + المياه الممغنطة وذلك خلال موسمي الشتاء على التوالي.